

Recent results from the LHC

From ATLAS and CMS

Alex Martyniuk (UCL)
on behalf of the ATLAS and CMS collaborations

Lake Louise Winter Institute
March 6th, 2020



Recent results from the LHC

From ATLAS and CMS


Alex Martyniuk (UCL)
on behalf of the ATLAS and CMS collaborations

RAL (remotely)
April 15th, 2020

The General Purpose Detectors

- This talk will focus on the results from ATLAS* and CMS





Differences

- Detector **technology** choices
- **B-field configuration**: Solenoid vs Solenoid+Toroid
- **Size/weight** (though both are colossal!)

Similarities

- **Cylindrical** detectors: barrel & end-caps
- **Concentric** detectors: Tracking, EM → had-calorimetry, muon chambers
- Close to 4π solid angle coverage
- Hardware/software **combined** trigger systems

CMS DETECTOR

Total weight : 14,000 tonnes
 Overall diameter : 15.0 m
 Overall length : 28.7 m
 Magnetic field : 3.8 T

STEEL RETURN YOKE

12,000 tonnes

SILICON TRACKERS

Pixel: (100x150 μm^2) - 10m² - 60M channels
 Microstrip: (10x100 μm^2) - 20m² - 64M channels

SUPERCONDUCTING SOLENOID

Niobium titanium coil carrying ~ 16,000A

MUON CHAMBERS

Barrel: 250 Drift Tubes, 400 Resonance Plate Chambers
 Endcaps: 400 Cathode Strips, 452 Resonance Plate Chambers

FRESHOWER

Silicon strips - 10x4" - 117,000 channels

FORWARD CALORIMETER


Steel + Quartz fibres - 2,000 Channels



CRYSTAL ELECTROMAGNETIC CALORIMETER (ECAL)

~76,000 scintillating PbWO₄ crystals



HADRON CALORIMETER (HCAL)

Iron + Plastic scintillator ~7,000 channels







- The LHC had an **excellent** Run 2, delivering $\approx 160 \text{ fb}^{-1}$ to both experiments!
- Both experiments recorded data with **superb** overall Run 2 data taking efficiency: ATLAS 94.2%, CMS 92.3%
 - This was achieved despite **challenging** pile-up conditions with $\langle \mu \rangle > 35$: i.e. on average 35 **simultaneous** $p - p$ collisions per bunch crossing!
- The LHC is very **versatile** machine, delivering in **special** runs throughout Run 2: $Pb - Pb$, $p - Pb$, $Xe - Xe$ and low pileup $p - p$ data to both experiments





- The LHC had an **excellent** Run 2, delivering $\approx 160 \text{ fb}^{-1}$ to both experiments!
- Both experiments recorded data with **superb** overall Run 2 data taking efficiency: ATLAS 94.2%, CMS 92.3%
 - This was achieved despite **challenging** pile-up conditions with $\langle \mu \rangle > 35$: i.e. on average 35 **simultaneous** $p - p$ collisions per bunch crossing!
- The LHC is very **versatile** machine, delivering in **special** runs throughout Run 2: $Pb - Pb$, $p - Pb$, $Xe - Xe$ and low pileup $p - p$ data to both experiments



- The LHC had an **excellent** Run 2, delivering $\approx 160 \text{ fb}^{-1}$ to both experiments!
- Both experiments recorded data with **superb** overall Run 2 data taking efficiency: ATLAS 94.2%, CMS 92.3%
 - This was achieved despite **challenging** pile-up conditions with $\langle \mu \rangle > 35$: i.e. on average 35 **simultaneous** $p - p$ collisions per bunch crossing!
- The LHC is very **versatile** machine, delivering in **special** runs throughout Run 2: $Pb - Pb$, $p - Pb$, $Xe - Xe$ and low pileup $p - p$ data to both experiments

- **Both** experiments have **excellent** reconstruction and calibration performance, even in the **harsh** pileup conditions of Run 2
- **Continuous** improvements seen in the calibrations, understandings of efficiencies, systematic uncertainties etc. over a **wide** p_T range
 - A **better** understanding of the detectors along with **data-driven** and **machine learning** techniques mean that object calibrations and efficiencies are often now **better** even in the harsher environment of Run 2




What is in this data?

- The LHC is an **EVERYTHING** factory (with additional background collisions)
 - Assuming 140 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$ both detectors have seen...

Particle	# produced
----------	------------

H boson	≈ 8 million
t quark	≈ 280 million
Z boson	≈ 8 billion
W boson	≈ 26 billion
b quark	≈ 160 trillion




- These datasets give both experiments **broad** physics programme potential
 - **High-precision** SM measurements, including Higgs properties
 - Detection of extremely **rare** processes
 - Exploration of **new** kinematic regimes for potential **new physics** signals!

What is in this data?

- The LHC is an **EVERYTHING** factory (with additional background collisions)
 - Assuming 140 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$ both detectors have seen...

Particle	# produced
----------	------------

H boson	≈ 8 million
t quark	≈ 280 million
Z boson	≈ 8 billion
W boson	≈ 26 billion
b quark	≈ 160 trillion



- These datasets give both experiments **broad** physics programme potential
 - **High-precision** SM measurements, including Higgs properties
 - Detection of extremely **rare** processes
 - Exploration of **new** kinematic regimes for potential **new physics** signals!





Image from [Symmetry Magazine](#)

Remarkable agreement



Standard Model Production Cross Section Measurements

Status: November 2019



The amazing **predictive** power of the Standard Model is **confirmed** by both experiments (**CMS versions**)


Agreement spans over **14 orders of magnitude**, with measurements/evidence of diverse and extremely rare SM processes

Improvements in theoretical calculations to NNLO level **complement** these results

Remarkable agreement

March 2019

CMS Preliminary




The amazing **predictive** power of the Standard Model is **confirmed** by both experiments (**CMS versions**)

Agreement spans over **14 orders of magnitude**, with measurements/evidence of diverse and extremely rare SM processes

Improvements in theoretical calculations to NNLO level **complement** these results

Remarkable agreement



Status: November 2019

- \square $pp \rightarrow X$
7 TeV, $20 \mu\text{b}^{-1}$, Nat. Commun. 2, 463 (2011)
- \square 8 TeV, $500 \mu\text{b}^{-1}$, Phys. Lett. B 761 158 (2016)
- \square 13 TeV, $60 \mu\text{b}^{-1}$, Phys. Rev. Lett. 117 182002 (2016)
- \triangle $pp \rightarrow W$ \square $pp \rightarrow Z/\gamma^*$
2.76 TeV, 4pb^{-1} , arXiv:1907.03567 (for Z/W)
- \square 5 TeV, 25pb^{-1} , Eur. Phys. J. C 79 (2019) 128 (for Z/W)
- \square 7 TeV, 4.6fb^{-1} , Eur. Phys. J. C 77 (2017) 367 (for Z/W)
- \square 8 TeV, 20.2fb^{-1} , JHEP 02, 117 (2017) (for Z)
- \square 8 TeV, 20.2fb^{-1} , Eur. Phys. J. C 79 (2019) 760 (for W)
- \square 13 TeV, 81pb^{-1} , PLB 759 (2016) 601 (for W)
- \square 13 TeV, 3.2fb^{-1} , JHEP 02, 117 (2017) (for Z)
- \square $pp \rightarrow t\bar{t}$
7 TeV, 4.6fb^{-1} , Eur. Phys. J. C 74:3109 (2014)
- \square 8 TeV, 20.3fb^{-1} , Eur. Phys. J. C 74:3109 (2014)
- \square 13 TeV, 3.2fb^{-1} , Phys. Lett. B 761 (2016)
- \square $pp \rightarrow tq$
7 TeV, 4.6fb^{-1} , PRD 90, 112006 (2014)
- \square 8 TeV, 20.3fb^{-1} , Eur. Phys. J. C 77 (2017) 531
- \square 13 TeV, 3.2fb^{-1} , JHEP 1704 (2017) 086
- \square $pp \rightarrow H$
7 TeV, 4.5fb^{-1} , Eur. Phys. J. C 76 (2016) 6
- \square 8 TeV, 20.3fb^{-1} , Eur. Phys. J. C 76 (2016) 6
- \square 13 TeV, 36.1fb^{-1} , Phys. Lett. B 786 (2018) 114
- \square $pp \rightarrow WW$
7 TeV, 4.6fb^{-1} , PRD 87, 112001 (2013)
- \square 8 TeV, 20.3fb^{-1} , JHEP 09 029 (2016)
- \square 13 TeV, 36.1fb^{-1} , arXiv:1905.04242
- \square $pp \rightarrow WZ$
7 TeV, 4.6fb^{-1} , Eur. Phys. J. C (2012) 72:2173
- \square 8 TeV, 20.3fb^{-1} , PRD 93, 092004 (2016)
- \square 13 TeV, 36.1fb^{-1} , arXiv:1902.05759
- \square $pp \rightarrow ZZ$
7 TeV, 4.6fb^{-1} , JHEP 03, 128 (2013)
- \square 8 TeV, 20.3fb^{-1} , JHEP 01, 099 (2017)
- \square 13 TeV, 36.1fb^{-1} , Phys. Rev. D 97 (2018) 032005

The amazing predictive power of the Standard Model is confirmed by both experiments (CMS versions)

Agreement spans over 14 orders of magnitude, with measurements/evidence of diverse and extremely rare SM processes

Improvements in theoretical calculations to NNLO level complement these results




Image from [Symmetry Magazine](#)


- The **discovery** of the Higgs boson ($m_H = 125.10$ GeV) in 2012 by ATLAS and CMS **together** fulfilled one of the **main aims** of the LHC: Identifying a **mass generation** mechanism for the SM
- Its discovery showed us that some form of the **Brout-Englert-Higgs*** mechanism is **realised** in nature!
- It has given us **access** to a new sector of the **SM Lagrangian** with new lines of enquiry to follow:
 - **Yukawa couplings**, a new type of interaction to investigate
 - Gauge–scalar boson **interactions**
 - The **parameters** of the Higgs potential, and its self coupling




*Please insert your own preferred naming convention for the mechanism

- In **Run 1** ATLAS and CMS observed the gg fusion and VBF production modes
- In **Run 2** we now have observed the Higgs-strahlung and ttH productions modes!


$ggF \sim 49\text{fb}@13\text{TeV}$




$VBF \sim 3.8\text{pb}@13\text{TeV}$




Higgs-strahlung



$bb/ttH \sim 0.5\text{pb}@13\text{TeV}$




- The **discovery** channels ($\gamma\gamma$, ZZ , WW) **dominated** Run 1 results
- In Run 2 both experiments are **digging** into the more **challenging** decay modes



Higgs Progress: Discovery → Measurement


Run 1: ATLAS & CMS





$$H \rightarrow \gamma\gamma$$

- Significant progress has been made in the discovery channels since 2012
- We have collected **thousands** of Higgs bosons candidate events with which to perform **differential** measurements
- Can really start to dig down into the **properties** and **couplings** of this new scalar boson

Run 2: ATLAS & CMS




Run 1: ATLAS & CMS





$$H \rightarrow WW$$

- Significant progress has been made in the discovery channels since 2012
- We have collected **thousands** of Higgs bosons candidate events with which to perform **differential** measurements
- Can really start to dig down into the **properties** and **couplings** of this new scalar boson

Run 2: ATLAS & CMS





Run 1: ATLAS & CMS



$$H \rightarrow ZZ$$

- **Significant** progress has been made in the discovery channels since 2012
- We have collected **thousands** of Higgs bosons candidate events with which to perform **differential** measurements
- Can really start to dig down into the **properties** and **couplings** of this new scalar boson





Run 2: ATLAS & CMS



Differential Higgs Measurements

ATLAS-CONF-2019-029 & ATLAS-CONF-2019-032 &
HIG-17-028 & HIG-19-002





- Statistical **combinations** of the large Higgs samples allows both experiments to provide **total** and **differential** cross-section results
- **Deviations** from the SM expectations in these measurements and in [Higgs couplings](#) could point us towards new physics
 - Could be subtle, → needs high precision



Differential Higgs Measurements

ATLAS-CONF-2019-029 & ATLAS-CONF-2019-032 & HIG-17-028 & HIG-19-002



- Statistical **combinations** of the large Higgs samples allows both experiments to provide **total** and **differential** cross-section results
- **Deviations** from the SM expectations in these measurements and in [Higgs couplings](#) could point us towards new physics
 - Could be subtle, → needs high precision



Two for one: $pp \rightarrow VH \rightarrow bb$

[HIGG-2018-04](#) & [HIG-18-016](#)

- **Observation** of the Higgs production/decay $pp \rightarrow VH \rightarrow bb$ **directly** confirms both the low cross section **Higgs-strahlung** production mode and the abundant $H \rightarrow bb$ **decay** mode
- Made possible by triggering on "**clean**" leptonic V decays to reduce the **multi-jet** background
- An **incredible** achievement as the $H \rightarrow bb$ decay channel was considered by some a **lost cause** at a hadron collider





- Combined Run 1+2 $H \rightarrow bb$ significances
–ATLAS–
 $5.4(5.5)\sigma$
–CMS–
 $4.8(4.9)\sigma$
- Both agree with the SM signal strength

The rarest production mode: $t\bar{t}H$


[HIGG-2018-13](#) & [HIG-17-035](#)

- **Combinations** of decay modes from both experiments have previously **confirmed** the presence of the $pp \rightarrow t\bar{t}H$ production channel at, ATLAS: $6.3(5.1)\sigma$, CMS: $5.2(4.2)\sigma$
- A **superb** confirmation of a **rare** Higgs production mode, confirming the **tree level coupling** of top quarks to the Higgs



[HIGG-2018-13](#) & [HIG-17-035](#) & [ATLAS-CONF-2019-004](#)

- **Combinations** of decay modes from both experiments have previously **confirmed** the presence of the $pp \rightarrow t\bar{t}H$ production channel at, ATLAS: $6.3(5.1)\sigma$, CMS: $5.2(4.2)\sigma$
- A **superb** confirmation of a **rare** Higgs production mode, confirming the **tree level coupling** of top quarks to the Higgs






- ATLAS has now followed up with a **single channel** ($H \rightarrow \gamma\gamma$) observation of the $t\bar{t}H$ process at a significance of 4.9σ
- An **exceptionally** rare process with a measured,
 $\sigma_{t\bar{t}H} \times \mathcal{B}_{H \rightarrow \gamma\gamma} = 1.59^{+0.43}_{-0.39} \text{ fb}$

The Rarer Decay Modes: 2nd Gen.

ATLAS-CONF-2019-028 & HIG-18-031



- ATLAS and CMS are digging down toward the **rare** second generation Higgs decay modes
- Using **multivariate** techniques and new reconstruction techniques
- **Nothing** is seen (or expected to be) seen yet, but these modes are starting their **journeys** now ready for Run 3/4
 - ATLAS $H \rightarrow \mu\mu$: $\mu = 1.7$ & CMS $H \rightarrow cc$: $\mu = 70$ @ 95%CL




The "Future": Di-Higgs production

[HDBS-2018-18](#) & [HDBS-18-33](#) & [ATLAS-CONF-2019-049](#) & [HIG-17-030](#) & [HIG-18-013](#)

- HH production can be searched for through **VBF** production modes





- Can be used to explore the Higgs **self coupling** λ , probing the **Higgs potential**
- Also useful to search for new **heavy resonances**



[HIGG-2018-57](#) & [HIG-19-005](#)

- The ATLAS and CMS Higgs programs are **switching** from discovery to measurements **quickly**
- Will be able to **compare** the results to the SM expectations with greater and greater **precision**
- The Higgs sector touches upon many **questions**: naturalness, vacuum stability, flavour...
- We will be **poking** this field for a long time to come










Image from [Symmetry Magazine](#)

ATLAS:1911.04813 CMS:1909.0413






- **High** precision (at percent level) measurements of **differential $Z\gamma$** cross-sections from **both** experiments
- In general **good** agreement seen with **NLO/NNLO** predictions
- Provides **high-statistics** probes of the electroweak gauge sector for the community to use




SMP-18-007 (ATLAS: [arXiv:1910.09503](https://arxiv.org/abs/1910.09503))

- **Evidence** of the electroweak vector-boson scattering process $pp \rightarrow Z\gamma jj$ **directly** probes the EWK SM gauge structure
- Selection **reduces** contribution from **strong** production
- Signal **extracted** from 2D fit to m_{jj} and $\Delta\eta_{jj}$





- Obs (exp) significance at $3.9 (5.2)\sigma$ with a cross-section of $3.20 \pm 1.15 \text{ fb}^{-1}$
- Additionally, places **stringent** limits on anomalous quartic gauge couplings

ATLAS-CONF-2019-033






- **Observation** of the electroweak vector-boson scattering process $pp \rightarrow ZZ_{jj}$ is a milestone in observing one of the **rarest processes** of the EW sector
- Selected from other QCD/EW diboson processes by a **Boosted Decision Tree**

- Observed at 5.5σ with a cross-section of 0.82 fb
 - One of the **rarest SM** process observed so far at the LHC
- CMS has plans ([FTR-18-014](#)) to measure the **longitudinal** component with 3 ab^{-1}



- **Evidence** of double parton scattering in the $pp \rightarrow W^\pm W^\pm$ channel is a **first** for DPS at the LHC
 - Could become a background to new physics with longitudinal correlations at HL-LHC, as well as other diboson processes
- **Multivariate** classifiers used to **discriminate** DPS events from other diboson backgrounds
- **Evidence** at 3.9σ with a cross-section of 1.41 pb^{-1}








Image from [Symmetry Magazine](#)

TOP-15-018

- Top quarks are **predominantly** produced in pairs in the SM
- **Anomalous** production modes can be searched for by studying the the **angular** distribution of the produced $t\bar{t}$ pairs



- I.e. these anomalies would **impact**,


$$c^* = \cos \theta^*$$

- This can be **quantified** by using the forward/backwards asymmetry,



$$A_{FB} = \frac{\sigma(c^* > 0) - \sigma(c^* < 0)}{\sigma(c^* > 0) + \sigma(c^* < 0)} \quad (A_{FB}^{SM} = 0.095)$$

- Trickier than at the Tevatron as gg fusion production is dominant at the LHC which produces no A_{FB} , need to extract $q\bar{q}$ contribution
- Can also measure the anomalous **chromoelectric** (\hat{a}_t) and **chromomagnetic** ($\hat{\mu}_t$) dipole moments ($\hat{a}_t^{SM} = \hat{\mu}_t^{SM} = 0$)



- Record **differential** cross sections in a series of channels **defined** by decay topology, lepton charge and flavour
- A linear **combination** of 3D MC templates is fitted to this data to **independently** extract
 - $A_{FB}^{(1)} = 0.048_{-0.087}^{+0.095}(\text{stat})_{-0.029}^{+0.020}(\text{syst})$
 - $\hat{\mu}_t = -0.024_{-0.009}^{+0.013}(\text{stat})_{-0.011}^{+0.016}(\text{syst})$
 - $|\hat{d}_t| < 0.03$ 95% CL






- Top quarks decay **before** their spins can be flipped by the strong interaction, passing this information to their **decay products**
- The initial $t\bar{t}$ spin states at the LHC depend on the **production mode** ($q\bar{q}$ annihilation/ gg fusion), \therefore a **different** measurement at Tevatron c.f. LHC
- Measurements by ATLAS and CMS generally agree with fixed order predictions (ATLAS has some tension with the NNLO prediction)
- ATLAS also has some tension with one NLO+PS prediction (POWHEG+PYTHIA8)



- The decay of $B_S^0 \rightarrow \mu^+\mu^-$ is **observed** by CMS with a branching fraction of $\mathcal{B}(B_S^0 \rightarrow \mu^+\mu^-) = 2.9_{-0.6}^{+0.7} \times 10^{-9}$ with a significance of 5.6 σ
 - **No significant excess** is seen for the $B^0 \rightarrow \mu^+\mu^-$, upper limits set at $\mathcal{B}(B^0 \rightarrow \mu^+\mu^-) = 3.6 \times 10^{-10}$
- The results are **consistent** with the SM expectation, and provide a significant **constraint** on BSM models which could enhance this channel



- A **precise** measurement of transverse momentum and ϕ_η^* (a measure of the lepton's scattering angle w.r.t. the beam) in **Drell-Yan** events
- High statistics measurements of standard candle processes are **important** inputs to beyond the Standard Model searches
- Unfolded differential cross section provide information to improve the **modelling** of these channels









Image (kinda) from [Symmetry Magazine](#)

- **Gravity**... It's just not in there...
- **Dark Matter**... Astronomers/Cosmologists say that it is everywhere...
 - The SM looks blankly into the distance...
 - Neutrino mass not enough, and SM doesn't even include them anyway...
- **Neutrino masses**... Also missing...
 - We know they have mass, the SM says they don't...
 - No (observed) ν_R so no Yukawa coupling...
 - What is the correct way to stick that in?
- A complete list of sticking plasters, things lacking an explanation and omissions would be **quite long**...
 - The Higgs mass, mass hierarchies in general, vacuum stability of the universe, multiple generations of fermions...
- And so we go **searching**...



Exotics & SUSY: Searching high and low

ATLAS Exotics Searches* - 95% CL Upper Exclusion Limits

Status: May 2019

ATLAS Preliminary

$$\int \mathcal{L} dt = (3.2 - 139) \text{ fb}^{-1}$$

$$\sqrt{s} = 8, 13 \text{ TeV}$$


	Model	ℓ, γ	Jets [†]	E_T^{miss}	$[\mathcal{L} dt[\text{fb}^{-1}]$	Limit	Reference
Extra dimensions	ADD $G_{\mu\kappa} + g/q$	$0, e, \mu$	1-4 J	Yes	36.1	M_{Pl} 7.7 TeV	$n = 2$ 1711.03031
	ADD non-resonant $\gamma\gamma$	$2, \gamma$	-	-	36.7	M_s 8.6 TeV	$n = 3$ HLZ NLO 1707.04147
	ADD QBH	-	2 J	-	37.0	M_{th} 8.9 TeV	$n = 6$ 1703.09127
	ADD BH high Σ, Δ	$\geq 1, e, \mu$	≥ 2 J	-	3.2	M_{th} 8.2 TeV	$n = 6, M_{\text{Pl}} = 3 \text{ TeV, rot BH}$ 1606.02265
	ADD BH mid Σ, Δ	$\geq 1, e, \mu$	≥ 3 J	-	3.6	M_{th} 9.55 TeV	$n = 6, M_{\text{Pl}} = 3 \text{ TeV, rot BH}$ 1512.02586
	RS1 $G_{\mu\kappa} \rightarrow \gamma\gamma$	$2, \gamma$	-	-	36.7	$G_{\mu\kappa}$ mass 4.1 TeV	$k/\bar{M}_{\text{Pl}} = 0.1$ 1707.04147
	Bulk RS $G_{\mu\kappa} \rightarrow WW/ZZ$	multi-channel	-	-	36.1	$G_{\mu\kappa}$ mass 2.3 TeV	$k/\bar{M}_{\text{Pl}} = 1.0$ 1808.02380
	Bulk RS $G_{\mu\kappa} \rightarrow WW \rightarrow q\bar{q}q\bar{q}$	$0, e, \mu$	2 J	-	139	$G_{\mu\kappa}$ mass 1.6 TeV	$k/\bar{M}_{\text{Pl}} = 1.0$ ATLAS-COFT-2019-003
	Bulk RS $G_{\mu\kappa} \rightarrow t\bar{t}$	$1, e, \mu$	$\geq 1, b, \geq 1J/2J$	Yes	36.1	$G_{\mu\kappa}$ mass 3.8 TeV	$\Gamma/m = 15\%$ 1804.10623
	2UED / RPP	$1, e, \mu$	$\geq 2, b, \geq 3J$	Yes	36.1	KK mass 1.8 TeV	Tier (1, 1), $\mathcal{R}(A^{(1,3)} \rightarrow t\bar{t}) = 1$ 1803.09678
Gauge bosons	SSM $Z' \rightarrow \ell\ell$	$2, e, \mu$	-	-	139	Z' mass 5.1 TeV	1903.06248
	SSM $Z' \rightarrow \tau\tau$	$2, \tau$	-	-	36.1	Z' mass 2.42 TeV	1709.07242
	Leptophobic $Z' \rightarrow b\bar{b}$	-	2 b	-	36.1	Z' mass 2.1 TeV	1805.09299
	Leptophobic $Z' \rightarrow t\bar{t}$	$1, e, \mu$	$\geq 1, b, \geq 1J/2J$	Yes	36.1	Z' mass 3.0 TeV	1804.10623
	SSM $W' \rightarrow \ell\nu$	$1, e, \mu$	-	-	139	W' mass 3.7 TeV	$\Gamma/m = 1\%$ CERN-EP-2019-100
	SSM $W' \rightarrow \nu\tau$	$1, \tau$	-	-	36.1	W' mass 6.0 TeV	1801.96992
	HVT $V' \rightarrow WZ \rightarrow q\bar{q}q\bar{q}$ model B	$0, e, \mu$	2 J	-	139	V' mass 3.7 TeV	$g_V = 3$ ATLAS-COFT-2019-003
	HVT $V' \rightarrow WH/ZH$ model B	multi-channel	-	-	36.1	V' mass 2.93 TeV	$g_V = 3$ 1712.00518
	LRSM $W_R \rightarrow t\bar{b}$	multi-channel	-	-	36.1	W_R mass 3.25 TeV	1807.10473
	LRSM $W_R \rightarrow \mu N_R$	$2, \mu$	1 J	-	80	W_R mass 5.0 TeV	$m(N_R) = 0.5 \text{ TeV, } g_L = g_R$ 1904.12679
CI	CI $q\bar{q}q\bar{q}$	-	2 J	-	37.0	A 21.8 TeV η_{CI}	1703.09127
	CI $\ell\ell q\bar{q}$	$2, e, \mu$	-	-	36.1	A 40.0 TeV η_{CI}	1707.02424
	CI $t\bar{t}t\bar{t}$	$\geq 1, e, \mu$	$\geq 1, b, \geq 1J$	Yes	36.1	A 2.57 TeV $ \zeta_{\text{CI}} = 4\pi$	1811.02305
DM	Axial-vector mediator (Dirac DM)	$0, e, \mu$	1-4 J	Yes	36.1	μ_{had} 1.55 TeV	$g_{\text{had}} = 0.25, g_{\text{le}} = 1.0, m(\chi) = 1 \text{ GeV}$ 1711.03031
	Colored scalar mediator (Dirac DM)	$0, e, \mu$	1-4 J	Yes	36.1	μ_{had} 1.67 TeV	$g_{\text{le}} = 1.0, m(\chi) = 1 \text{ GeV}$ 1711.03031
	$W_{\text{V}1,2}$ EFT (Dirac DM)	$0, e, \mu$	1 J, $\leq 1J$	Yes	3.2	M_{th} 700 GeV	$m(\chi) < 150 \text{ GeV}$ 1608.02372
	Scalar reson. $\phi \rightarrow t\bar{t}$ (Dirac DM)	$0-1, e, \mu$	1 b, 0-1 J	Yes	36.1	M_{th} 3.4 TeV	$\gamma = 0.4, a = 0.2, m(\chi) = 10 \text{ GeV}$ 1812.09743
LQ	Scalar LQ 1 st gen	$1, 2, e, \mu$	≥ 2 J	Yes	36.1	LQ mass 1.4 TeV	$\beta = 1$ 1902.00377
	Scalar LQ 2 nd gen	$1, 2, \mu$	≥ 2 J	Yes	36.1	LQ mass 1.56 TeV	$\beta = 1$ 1902.00377
	Scalar LQ 3 rd gen	$2, \tau$	2 b	-	36.1	LQ mass 1.03 TeV	$\mathcal{R}(LQ_{\text{c}}^{\pm} \rightarrow b\bar{r}) = 1$ 1902.08103
	Scalar LQ 3 rd gen	$0-1, e, \mu$	2 b	Yes	36.1	LQ mass 970 GeV	$\mathcal{R}(LQ_{\text{c}}^{\pm} \rightarrow t\bar{r}) = 0$ 1902.08103
	VLO $TT \rightarrow HZ/Zt/Wb + X$	multi-channel	-	-	36.1	T mass 1.37 TeV	SU(2) doublet 1808.02343
VLO $BB \rightarrow Wt/Zb + X$	multi-channel	-	-	36.1	B mass 1.34 TeV	SU(2) doublet 1808.02343	
VLO $T_{3,1/3} T_{3,1/3} \rightarrow Wt + X$	$2(S_{\text{SU}}) \geq 3, e, \mu$	$\geq 1, b, \geq 1J$	Yes	36.1	$T_{3,1/3}$ mass 1.64 TeV	$\mathcal{R}(T_{3,1/3} \rightarrow Wt) = 1$ 1807.11883	
VLO $Y \rightarrow Wb + X$	$1, e, \mu$	$\geq 1, b, \geq 1J$	Yes	36.1	Y mass 1.85 TeV	$\mathcal{R}(Y \rightarrow Wb) = 1, c_{\text{th}}(Wb) = 1$ 1812.07343	
VLO $B \rightarrow Hb + X$	$0, e, \mu, 2, \gamma$	$\geq 1, b, \geq 1J$	Yes	79.8	B mass 1.21 TeV	$\kappa_{\text{th}} = 0.5$ ATLAS-COFT-2018-024	
VLO $QQ \rightarrow WbWq$	$1, e, \mu$	$\geq 4J$	Yes	20.3	Q mass 690 GeV	1509.04261	
Excited fermions	Excited quark $q^* \rightarrow q\bar{q}$	-	2 J	-	139	q^* mass 6.7 TeV	only u^* and d^* , $\Lambda = m(q^*)$ ATLAS-COFT-2019-007
	Excited quark $q^* \rightarrow q\gamma$	1 γ	1 J	-	36.7	q^* mass 5.3 TeV	1709.10440
	Excited quark $q^* \rightarrow qg$	-	1 b, 1 J	-	36.1	q^* mass 2.6 TeV	1805.09299
	Excited lepton ℓ^*	$3, e, \mu$	-	-	20.3	ℓ^* mass 3.0 TeV	$\Lambda = 3.0 \text{ TeV}$ 1411.2921
	Excited lepton ν^*	$3, e, \mu, \tau$	-	-	20.3	ν^* mass 1.6 TeV	$\Lambda = 1.6 \text{ TeV}$ 1411.2921
Other	Type III Seesaw	$1, e, \mu$	≥ 2 J	Yes	79.8	N^c mass 560 GeV	ATLAS-COFT-2019-020
	LRSM Majorana	$2, \mu$	2 J	-	36.1	N_{th} mass 3.2 TeV	$m(W_{\text{C}}) = 4.1 \text{ TeV, } g_L = g_R$ 1809.11105
	Higgs triplet $H^{\pm\pm} \rightarrow \ell\ell$	$2, 3, 4, e, \mu$ (SS)	-	-	36.1	$H^{\pm\pm}$ mass 870 GeV	DY production, $\mathcal{R}(H^{\pm\pm} \rightarrow t\bar{t}) = 1$ 1710.09748
	Higgs triplet $H^{\pm\pm} \rightarrow t\bar{t}$	$3, e, \mu, \tau$	-	-	20.3	$H^{\pm\pm}$ mass 400 GeV	DY production, $\mathcal{R}(H^{\pm\pm} \rightarrow t\bar{t}) = 1$ 1411.2921
	Multi-charged particles	-	-	-	36.1	multi-charged particle mass 1.22 TeV	DY production, $ \zeta = 5\pi$ 1812.03673
	Magnetic monopoles	-	-	-	34.4	monopole mass 2.37 TeV	DY production, $ \zeta = 1g_2, \text{spin } 1/2$ 1905.10130
		$\sqrt{s} = 8 \text{ TeV}$	$\sqrt{s} = 13 \text{ TeV}$ partial data	$\sqrt{s} = 13 \text{ TeV}$ full data			

*Only a selection of the available mass limits on new states or phenomena is shown.

†Small-radius (large-radius) jets are denoted by the letter (L)(R).

Overview of CMS EXO results

36 fb⁻¹ (13 TeV)



Selection of observed exclusion limits at 95% C.L. (theory uncertainties are not included).

January 2019

Exotics & SUSY: Searching high and low



ATLAS SUSY Searches* - 95% CL Lower Limits

July 2019

ATLAS Preliminary

$\sqrt{s} = 13 \text{ TeV}$


Model	Signature	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Mass limit	Reference			
Inclusive Searches	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow \text{gl}$ ⁰	0 e, μ mono-jet	36.1 36.1	\tilde{g} [9K, 8k Degrad] \tilde{g} [1k, 8k Degrad]	$m(\tilde{g}) > 100 \text{ GeV}$ $m(\tilde{g}) = m(\tilde{t}) > 5 \text{ GeV}$		
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow \text{gg}$ ⁰	0 e, μ 2-6 jets	$E_{\text{miss}}^{\text{min}}$ $E_{\text{miss}}^{\text{max}}$	36.1 36.1	\tilde{g} \tilde{g}	$m(\tilde{g}) > 200 \text{ GeV}$ $m(\tilde{g}) > 900 \text{ GeV}$	
	$3e, \mu$ $\tilde{g}\tilde{g}, \tilde{g} \rightarrow \text{gg}(CR)\tilde{g}$ ⁰	4 jets e, μ, ν	36.1 36.1	Forbidden	1.55 2.0	$m(\tilde{g}) > 400 \text{ GeV}$ $m(\tilde{g}) > 300 \text{ GeV}$	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow \text{gg}WZ\tilde{g}$ ⁰	0 e, μ 7-11 jets	$E_{\text{miss}}^{\text{min}}$ 139	\tilde{g}	1.2 1.8	$m(\tilde{g}) > 400 \text{ GeV}$ $m(\tilde{g}) > 200 \text{ GeV}$	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow \text{gg}$ ⁰	0-1 e, μ 3 b	$E_{\text{miss}}^{\text{min}}$ 79.8	\tilde{g}	1.15	$m(\tilde{g}) > 200 \text{ GeV}$	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow \text{gg}$ ⁰	0-1 e, μ 6 jets	$E_{\text{miss}}^{\text{min}}$ 139	\tilde{g}	1.25	$m(\tilde{g}) > 300 \text{ GeV}$	
	3 rd gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow \text{bb}\tilde{t}_1^0\tilde{t}_1^0$	Multiple Multiple	36.1 36.1 139	Forbidden Forbidden	0.9 0.58-0.82 0.74	$m(\tilde{t}) > 300 \text{ GeV}, \text{BR}(\tilde{t}_1^0 \rightarrow \tilde{b}_1\tilde{t}_1^0) = 1$ $m(\tilde{t}) > 300 \text{ GeV}, \text{BR}(\tilde{t}_1^0 \rightarrow \tilde{b}_1\tilde{t}_1^0) = 0.5$ $m(\tilde{t}) > 200 \text{ GeV}, m(\tilde{t}) > 300 \text{ GeV}, \text{BR}(\tilde{t}_1^0 \rightarrow \tilde{t}_1^0) = 1$
		$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow \text{bb}\tilde{t}_1^0 \rightarrow \text{bb}\tilde{t}_1^0$	0 e, μ 6 b	$E_{\text{miss}}^{\text{min}}$ 139	Forbidden	0.23-0.48	$\Delta m(\tilde{t}_1^0, \tilde{b}_1) > 130 \text{ GeV}, m(\tilde{t}_1^0) > 100 \text{ GeV}$ $\Delta m(\tilde{t}_1^0, \tilde{b}_1) > 130 \text{ GeV}, m(\tilde{t}_1^0) > 0 \text{ GeV}$
		$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{Wb}\tilde{b}_1^0$ or $\tilde{t}_1^0\tilde{t}_1^0$	0-2 e, μ 0-2 jets/1-2 b	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{t}_1^0	1.0	$m(\tilde{t}_1^0) > 1 \text{ GeV}$
		$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{Wb}\tilde{b}_1^0$	1 e, μ 3 jets/1 b	$E_{\text{miss}}^{\text{min}}$ 139	\tilde{t}_1^0	0.44-0.59	$m(\tilde{t}_1^0) > 600 \text{ GeV}$
$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{Wb}\tilde{b}_1^0, \tilde{t}_1^0 \rightarrow \text{t}\tilde{g}$		1 + 1 e, μ 2 jets/1 b	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{t}_1^0	1.15	$m(\tilde{t}_1^0) > 800 \text{ GeV}$	
$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{Wb}\tilde{b}_1^0, \tilde{t}_1^0 \rightarrow \text{t}\tilde{g}, \tilde{t}_1^0 \rightarrow \text{bb}\tilde{t}_1^0$		0 e, μ 2 c	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{t}_1^0	0.46 0.85	$m(\tilde{t}_1^0) > 0 \text{ GeV}$ $m(\tilde{t}_1^0) > 300 \text{ GeV}$	
$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{Wb}\tilde{b}_1^0$		0 e, μ mono-jet	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{t}_1^0	0.43	$m(\tilde{t}_1^0) > 300 \text{ GeV}, m(\tilde{t}_1^0) > 5 \text{ GeV}$	
$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \tilde{t}_1^0 + h$		1-2 e, μ 4 b	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{t}_1^0	0.32-0.88	$m(\tilde{t}_1^0) > 0 \text{ GeV}, m(\tilde{t}_1^0) > 180 \text{ GeV}$	
$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \tilde{t}_1^0 + Z$		3 e, μ 1 b	$E_{\text{miss}}^{\text{min}}$ 139	Forbidden	0.86	$m(\tilde{t}_1^0) > 360 \text{ GeV}, m(\tilde{t}_1^0) > 40 \text{ GeV}$	
EW direct		$\tilde{t}_1^0\tilde{t}_1^0$ via WZ	2-3 e, μ e, μ, ν	$E_{\text{miss}}^{\text{min}}$ 139	$\tilde{t}_1^0\tilde{t}_1^0$	0.205	$m(\tilde{t}_1^0) > 190, 0290$ $m(\tilde{t}_1^0) = m(\tilde{t}_1^0) > 5 \text{ GeV}$
	$\tilde{t}_1^0\tilde{t}_1^0$ via WW	2 e, μ $E_{\text{miss}}^{\text{min}}$	139	\tilde{t}_1^0	0.42	$m(\tilde{t}_1^0) > 0$	
	$\tilde{t}_1^0\tilde{t}_1^0$ via Wb	0-1 e, μ 2 b/2 ν	$E_{\text{miss}}^{\text{min}}$ 139	$\tilde{t}_1^0\tilde{t}_1^0$	Forbidden	0.74	$m(\tilde{t}_1^0) > 70 \text{ GeV}$
	$\tilde{t}_1^0\tilde{t}_1^0$ via $\tilde{t}_1^0\tilde{t}_1^0$	2 e, μ $E_{\text{miss}}^{\text{min}}$	139	\tilde{t}_1^0	1.0	$m(\tilde{t}_1^0) > 0.5m(\tilde{t}_1^0) + m(\tilde{t}_1^0)$	
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{t}\tilde{g}$	2 e, μ 0 jets	$E_{\text{miss}}^{\text{min}}$ 139	\tilde{t}_1^0	0.16-0.3 0.12-0.39	$m(\tilde{t}_1^0) > 0$	
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{t}\tilde{g}$	2 e, μ ≥ 1 b	$E_{\text{miss}}^{\text{min}}$ 139	\tilde{t}_1^0	0.256	$m(\tilde{t}_1^0) > 10 \text{ GeV}$	
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{t}\tilde{g}$	2 e, μ ≥ 3 b	$E_{\text{miss}}^{\text{min}}$ 139	\tilde{t}_1^0	0.13-0.23	$m(\tilde{t}_1^0) > 10 \text{ GeV}$	
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{t}\tilde{g}$	4 e, μ 0 jets	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{t}_1^0	0.3	$\text{BR}(\tilde{t}_1^0 \rightarrow \tilde{t}_1^0) > 1$ $\text{BR}(\tilde{t}_1^0 \rightarrow \tilde{t}_1^0) > 1$	
	Long-lived particles	Direct $\tilde{t}_1^0\tilde{t}_1^0$ prod., long-lived \tilde{t}_1^0	Disapp. tik 1 jet	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{t}_1^0	0.15	0.46
		Stable \tilde{g} R-hadron Metastable \tilde{g} R-hadron, $\tilde{g} \rightarrow \text{gg}\tilde{t}_1^0$	Multiple Multiple	36.1 36.1	\tilde{g}	2.0 2.05, 2.4	$m(\tilde{g}) > 100 \text{ GeV}$
RPV	LFV $\tilde{p}\tilde{p} \rightarrow \tilde{p}\tilde{p} + X, \tilde{p} \rightarrow \text{qq}\tilde{t}_1^0\tilde{t}_1^0$	e, μ, τ, ν 0 jets	$E_{\text{miss}}^{\text{min}}$ 36.1	\tilde{p}	1.9	$\tilde{p}_{\text{LSP}} = 0.11, A_{10,11,12} = 0.07$	
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{W}\tilde{W}\tilde{Z}\tilde{t}_1^0\tilde{t}_1^0$	4 e, μ 0 jets	$E_{\text{miss}}^{\text{min}}$ 36.1	$\tilde{t}_1^0\tilde{t}_1^0$	0.82	$m(\tilde{t}_1^0) > 100 \text{ GeV}$	
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{gg}\tilde{t}_1^0$	Multiple 4-5 large R-jets	36.1 36.1	\tilde{t}_1^0	1.33 1.3	1.9 2.0	$m(\tilde{t}_1^0) > 200 \text{ GeV}, 1100 \text{ GeV}$ $m(\tilde{t}_1^0) > 200 \text{ GeV}, 1100 \text{ GeV}$
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{gg}\tilde{t}_1^0$	Multiple 2 jets + 2 b	36.1 36.7	\tilde{t}_1^0	0.36 0.42	1.05	$m(\tilde{t}_1^0) > 200 \text{ GeV}, \text{bino-like}$
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{bb}\tilde{t}_1^0$	2 e, μ 2 b	36.1 36.7	\tilde{t}_1^0	0.42	0.61	$m(\tilde{t}_1^0) > 200 \text{ GeV}, \text{bino-like}$
	$\tilde{t}_1^0\tilde{t}_1^0, \tilde{t}_1^0 \rightarrow \text{gg}\tilde{t}_1^0$	1 μ DV	139 139	\tilde{t}_1^0	1.0	0.4-1.45 1.6	$\text{BR}(\tilde{t}_1^0 \rightarrow \text{t}\tilde{g}) > 20\%$ $\text{BR}(\tilde{t}_1^0 \rightarrow \text{t}\tilde{g}) > 100\%, \text{cos}\theta = 1$

*Only a selection of the available mass limits on new states or phenomena is shown. Many of the limits are based on simplified models, c.f. refs. for the assumptions made.

10⁻¹ 1 Mass scale [TeV]

Selected CMS SUSY Results* - SMS Interpretation

ICHEP '16 - Moriond '17



CMS Preliminary


$\sqrt{s} = 13\text{TeV}$

$L = 12.9 \text{ fb}^{-1}$ $L = 35.9 \text{ fb}^{-1}$

More up to date, but separated summaries can be found [here](#)

- Searches for exotic phenomena and SUSY partners push to extremes of phase space
- They also often turn the normal operation of the detectors upside down





- These **extremes** include...
- The **highest** invariant mass events
- **Low** mass particles
- **Compressed** spectra
- **Small** couplings
- **Long-lived** particles
- **Multi-charged** particles
- **Forbidden** decays
- **Complicated** decays



A $m_{jj} = 9.3$ TeV dijet event recorded by ATLAS





EXOT-2019-03 & EXO-19-012

- Dijet resonance searches probe the **highest** invariant mass events, $m_{jj} = 9.3$ TeV now for ATLAS, $m_{jj} = 8.2$ TeV in CMS
- This iteration from ATLAS is **expanding** on the inclusive search by requiring **additional** b -tagging selections to probe other models
- **No** excesses are seen in any of the spectra



[EXOT-2018-08](#) & [EXOT-2018-30](#) & [EXO-19-019](#) & [EXO-16-033](#)





- Dilepton resonance searches are **complimentary** searches to the dijet searches, exploring a **wide range** of BSM models
- Require **careful** treatment of SM backgrounds out to **high** invariant mass regions
- Again **no** excesses are seen in any of the spectra



"Low" mass resonances




EXO-19-018&EXO-18-012&EXOT-2018-05&EXOT-2016-20

- Can **access** lower mass resonances than triggers would usually allow by **recoiling** the system against ISR, or by running on **reduced** size trigger level data
- Multiple examples from **both** experiments, pushing searches into areas **previously** thought inaccessible to the LHC
- **No excesses** seen in any of these searches either, but they demonstrate the **ingenuity** of collaboration members






SUS-19-007

- A search for gluino pairs with a **spectacular fireworks display** of a final state of $t\bar{t}\bar{t} + p_T^{\text{miss}}$
- The variables M_J , the scalar sum of large-R jets, and M_T , transverse mass of leading lepton + p_T^{miss} , offer **strong handles** on these busy events
- Regions ($R_{1,2,3}$) **dominated** by background are used to estimate the background in the signal region (R_4)
- Data is able to **exclude** gluinos with masses below 2.15 TeV



- Searches also ongoing for **production** of electroweakinos, **close** to the EW mass scale, and in **compressed** spectra
- Push back into regions where the $m(\tilde{\chi}_2^0) \approx m(\tilde{\chi}_1^0) + m(Z)$ or $m(\tilde{\chi}_1^0) = m(\tilde{\chi}_2^0)$
- The "easier" search regions are longer term statistics driven games now









Image courtesy of [Heather Russell](#)

EXOT-2018-61




- Searches for LLPs require **novel** trigger and reconstruction techniques, often a complete **re-write**
- **New** reconstruction methods developed for displaced tracks in the ID and MS detectors
- **Unusual** backgrounds from material interactions, fake vertices or punch through jets
- Place **limits** on the $c\tau$ of the long lived particles



Long-Lived Particles: DNN Jet Tagging

EXO-19-011


- CMS has **developed** a novel tagger to identify jets originating from LLP events
- Uses a Deep Neural Network which **achieves** a tagging efficiency of 30-80% for gluinos with $1\text{ mm} \leq c\tau_0 \leq 10\text{ m}$
- Expect an **improvement** in limits over **previous** results by using this **novel** technique



We are not done yet...





LHC / HL-LHC Plan



...not by a **long** way

● Liquid Argon Calorimeter Electronics

- Aiming to improve the Level-1 calorimeter decisions for Run 3 and beyond
- Finer segmentation leading to enhanced jet rejection and pileup subtraction capabilities




● TDAQ upgrades


- Take advantage of finer segmentation in LAr electronics and improved muon trigger (NSW)

● Muons: New Small Wheel

- Replacing the inner muon stations in the endcaps
- Reduced muon fake rates, and maintain the same position resolution/efficiency for HL-LHC



- CMS is well advanced on its Phase-1 upgrades
 - Upgrades of the pixels, L1 Trigger system and replacement of some HCAL readout have already occurred
 - Final upgrades ongoing in LS2 including replacing the inner layer of the pixel barrel




● Hadronic Calorimeter Electronics

- Replace photosensors of hybrid photodetectors with silicon photomultipliers
- Will improve the readout to 5Gbps and increase the longitudinal segmentation of the detector

● Muons: **GEM GEI/I detectors**

- Technically already a Phase-2 upgrade, but going in now in LS2
- Installing Gas-Electron-Multiplier chambers which can operate in high-rate environments






LHC / HL-LHC Plan




...not by a **long** way

We are not done yet...





There is a lot of data incoming... **pessimistically** 3 ab^{-1}



There is a lot of data incoming... or **optimistically** $4++ \text{ ab}^{-1}$

That data will be at $\langle \mu \rangle = 200$





Upgraded Trigger and Data Acquisition System:

- L0: 1 MHz
- Improved High-Level Trigger

Electronics Upgrade :

- LAr Calorimeter
- Tile Calorimeter
- Muon system

New Inner Tracking Detector (all silicon tracker, up to $|\eta| = 4$)

New muon chambers
in the inner barrel region

Options:

- High granularity timing detector (forward region)
- High- η muon tagger

LI Trigger/HLT/DAQ

- LI 40 MHz in/750 kHz out with tracking for PF-like
- HLT 7.5 kHz out

Beam Radiation and Luminosity, Common Systems, Infrastructure

Calorimeter Endcap

- Si, Scint + SiPM in Pb-W-SS
- 3D shower imaging with precise timing

Tracker

- Si Strip Outer Tracker designed for LI Track Trigger
- Pixelated Inner Tracker extends coverage to $|\eta| < 3.8$

Barrel Calorimeters

- ECAL single crystal granularity in LI Trigger with precise timing for e/γ at 30 GeV
- ECAL and HCAL new back-end electronics


Muon Systems

- DT & CSC new FE/BE readout
- New GEM/RPC $1.6 < |\eta| < 2.4$
- Extended coverage to $|\eta| < 2.8$

$|\eta| < 3.0$



MIP Timing Detector

- 30 ps resolution
- Barrel: Crystals + SiPMs
- Endcap: LGADs



- Both of the upgrade programs are **major undertakings**
- Equivalent to (re-)building a **good fraction** of the detectors in each case, but while the collaboration is **still running** the existing system and producing physics results
- **Very challenging!**

- ATLAS & CMS are **ploughing** through their $\approx 140\text{fb}^{-1}$ Run 2 datasets
- **No** new physics seen, but **exciting** results none the less
 - **Precision** differential Higgs cross section measurements, **progress** in rare Higgs decays and constraints on the $VVHH$ system
 - **Observation** of the $WWjj$, $WZjj$, $ZZjj$ electroweak scattering processes
 - **Precision** measurements of SM processes, increasing our understanding and constraining backgrounds to new physics
 - **Powerful** searches continue for new physics exploring new signatures and new parameter space
- Both experiments are **preparing** for the challenges of Run 3 and beyond into HL-LHC
- We already have a **gold mine** of experimental data, soon* we will be **spoiled** with $10\times$ as much data!



*For a generous definition of soon

