



ALICE



UNIVERSITY OF  
LIVERPOOL

# Uncovering the microscopic structure of the quark-gluon plasma

Jaime Norman

University of Liverpool

Nuclear Physics Community Meeting Daresbury Lab

8th January 2026

# Many-body QCD and the quark-gluon plasma (QGP)





fig. H. Caines

- Phase transition at high temperature or density to deconfined state of quarks and gluons  
- **quark-gluon plasma (QGP)**
- Calculations on the lattice predicts smooth crossover at  $\sim 155$  MeV at low baryon density
- Created using **ultra-relativistic heavy-ion collisions**

→ by studying these collisions we study the bulk, high-temperature properties of QCD matter, and the emergent phenomena that may arise

# Where can we make a QGP?



- Collisions of protons (pp), gold-ions (Au-Au), plus many other ions (Al, Cu, Zr, U...)
- Centre-of-mass energy per nucleon-nucleon collision up to  $\sqrt{s_{NN}} = 200$  GeV (Au-Au)




- In addition to collisions of protons (pp), for one month a year collide lead-ions (Pb-Pb) and proton-lead (p-Pb) (plus Xe-Xe, this year O-O, Ne-Ne)
- Centre-of-mass energy per nucleon-nucleon collision up to  $\sqrt{s_{NN}} = 5.36$  TeV (Pb-Pb),  $\sqrt{s} = 13.6$  TeV (pp)

# Heavy-ion collisions at the LHC

## The ALICE detector

The dedicated heavy-ion experiment at the LHC



- **Charged particle tracking** performance over a wide momentum range ( $\sim 0.15 - 100$  GeV/c)
- **Particle identification** (TPC, TOF) and photon/neutral ID (EMCal, Photon spectrometer)
- **Muon spectrometer** at forward rapidities


**Major upgrade** for 2022-2026 runs (LHC Run 3)

- **x2 improvement in spatial precision**
- **>x100 increase in collected collisions w.r.t Runs 1 and 2**

# QGP (in a nutshell)

## Long-distance structure:

QGP is a strongly-coupled liquid  
(with very low viscosity)



P. Romatschke

$$\eta/s \sim 280$$


$$\eta/s \sim 0.12$$

- Lower bound from strongly-coupled gauge theory  
 $\sim 1 / 4\pi \sim 0.08$

**The 'perfect liquid'!**

## Short distance structure:

Free quarks and gluons? Complex bound states?  
degrees of freedom not yet established



**What is the (quasi-particle) structure of the QGP as a function of resolution scale?**

# Jets: Multi-scale 'probes' of the QGP


Evolution of hard parton (quark or gluon)

→ gluon radiation

- Experimentally measured as **collimated spray of hadrons**

Reconstruct jets


→ measure initiating parton



'Jet quenching' - partonic interactions in the QGP


- inelastic (medium-induced gluon emission) and elastic (collisional) processes over full parton shower

Jets provide unique probes of the QGP as a function of its resolution scale





# Designing a scattering experiment

Ernest Rutherford  
uncovered atomic structure




Can we uncover  
microscopic quark-gluon  
plasma structure?



# Designing a scattering experiment

Theory state of the art:



- **Strong-coupling limit** - probability of parton to obtain momentum  $k_T$  is Gaussian (exponential) distributed
- If medium probed at short enough distance scales - scatter off **weakly-interacting quasiparticle**  
~ 'Rutherford-Like' power-law distributed  $\sim 1/(k_T)^4$

F. D'eraimo, M. Lekaveckas, H. Liu, K. Rajagopal, *JHEP* 05 (2013) 031

F. D'eraimo, K. Rajagopal, Y. Yin *JHEP* 01 (2019)

- Radiative corrections lead to harder power law  $1/(k_T)^{4-2\beta}$   
- hard scatters more likely

P. Caucal, Y. Mehtar-Tani: *Phys.Rev.D* 106 (2022) 5, L051501

*JHEP* 09 (2022) 023

*Phys.Rev.D* 108 (2023) 1, 014008


→ search for low-energy, large-angle scatterings of jets or their constituents  
'Molière scattering'

# Designing a scattering experiment


Experiment state of the art: search for large-angle angular broadening of a dijet system

ALICE: Phys.Rev.Lett. 133 (2024) 2, 022301, Phys.Rev.C 110 (2024) 1, 014906

See also [JN CERN LHC seminar](#)



- Semi-inclusive measurement of jets recoiling from high- $p_T$  hadron
- Low energy jets ( $\sim 10$  GeV) accessed via background subtraction techniques




# Designing a scattering experiment


Experiment state of the art: search for large-angle angular broadening of a dijet system

ALICE: Phys.Rev.Lett. 133 (2024) 2, 022301, Phys.Rev.C 110 (2024) 1, 014906

See also JN CERN LHC seminar



- Semi-inclusive measurement of jets recoiling from high- $p_T$  hadron
- Low energy jets ( $\sim 10$  GeV) accessed via background subtraction techniques



→ **First observation of ‘jet acoplanarity’ - predicted in 80’s!**

Appel, 1986  
 Blaizot, McLerran, 1986

# Designing a scattering experiment



Experiment state of the art: search for large-angle angular broadening of a dijet system

ALICE: Phys.Rev.Lett. 133 (2024) 2, 022301, Phys.Rev.C 110 (2024) 1, 014906

See also [JN CERN LHC seminar](#)

Dependence on jet aperture (radius ' $R$ ') and model comparisons

→ primarily caused by '*shockwave*' (wake) in quark-gluon plasma rather than large-angle scatterings




→ **First observation of 'jet acoplanarity' - predicted in 80's!**

Appel, 1986  
 Blaizot, McLerran, 1986

# Designing a scattering experiment

Experiment state of the art: search for large-angle scattering of jet splittings

**Parton  
inside jet**




- **Idea:** Measure relative transverse momentum ( $k_T$ ) of jet splittings in the parton shower
- **In practice:** Sequential jet clustering/ declustering algorithms give access to parton shower history

0–10%, 30–50% Pb–Pb, pp  $\sqrt{s_{NN}} = 5.02$  TeV  
 Anti- $k_T$  ch-particle jets,  $R = 0.2$ ,  $|\eta_{jet}| < 0.7$   
 $60 < p_{T, ch jet} < 80$  GeV/c





ALICE: Phys. Rev. Lett. 135, 031901 (2025)

■ Hybrid w/ wake  
■ Hybrid w/ wake + Moliere  
■ JETSCAPEv3.5 AA22  
■ JEWEL (recoils)  
■ JEWEL (no recoils)



→ **No clear evidence of large-angle scattering**

# Next steps





- Can we **separate** the jet-induced ‘*shockwaves*’ from the ‘*scatters*’ ?
- Can we **define a precise probe** sensitive to in-medium scattering?
- How can we rigorously **validate models** of the plasma’s microscopic structure?


# Constraining interplay between wake and jets

Challenge: 'Wake' from jet-medium interactions may mask hard scatterings

**Next steps:** characterise wake effects via measurement of *jet internal structure*




Qin, Majumder, Song, Heinz,  
Phys. Rev. Lett. 103, 152303 (2009)



arxiv:2512.12715


Significant interest in its own right - shows how the quark-gluon plasma responds to excitation - constrain viscosity, equation of state...

# Constraining interplay between wake and jets





- Low-energy ‘broadened’ jets expected to have significant contribution from **thermalised particles** from the wake
- Studying internal structure of these jets will confirm thermal nature  
→ **new window into studying the QGP and its transport properties**
- Characterise thermal component  
→ **more precise window into ‘hard’ scattering within the QGP**

# Constraining interplay between wake and jets




- Low-energy ‘broadened’ jets expected to have significant contribution from **thermalised particles** from the wake
- Studying internal structure of these jets will confirm thermal nature  
→ **new window into studying the QGP and its transport properties**
- Characterise thermal component  
→ **more precise window into ‘hard’ scattering within the QGP**


Jet  
transverse  
energy  
distribution




Jet axes  
difference




# Constraining interplay between wake and jets




- Low-energy ‘broadened’ jets expected to have significant contribution from **thermalised particles** from the wake
- Studying internal structure of these jets will confirm thermal nature  
→ **new window into studying the QGP and its transport properties**
- Characterise thermal component  
→ **more precise window into ‘hard’ scattering within the QGP**



ALICE Run 3  
(uncorrected) data



Jet axes difference






# Measuring a precise scattering probe

**Challenge:** need precise handle on scattering probe  
(time evolution, initial energy...)

**Next steps:** precision measurement of  
*charm-hadron correlations and  
charm hadron-tagged jets*

heavy quarks (charm + beauty) only produced in initial stages of collision

→ **tangible, traceable probe of scatters**




F. Prino, R. Rapp, *J.Phys.G* 43 (2016) 9, 093002

L. Apolinario, Y.-J. Lee, and M. Winn, *Prog. Part. Nucl. Phys.* 127 (2022) 103990


# Measuring a precise scattering probe

State of the art: inclusive charm hadron production



ALI-PUB-561427

$$R_{AA}(p_T) = \frac{1}{\langle N_{coll} \rangle} \frac{dN_{AA}/dp_T}{dN_{pp}/dp_T}$$



Exploration of *new, multi-differential methods* to measure charm correlations, and to ‘tag’ charm jets ongoing

# Connecting data with theory




The tool for **comprehensive validation** of cutting-edge models with multi-messenger data

→ **does a model capture the essential physics?**

## Heavy-ion event generator

Framework for modelling all aspects of heavy-ion collisions



## Statistical toolkit

Bayesian inference for data-theory comparison

$$P(\theta | \text{data}) = \frac{P(\text{data} | \theta)P(\theta)}{P(\text{data})}$$

**Modular** - substitute in different models

**Interdisciplinary**

(experimentalists, theorists, statisticians)

**Computing intensive calculations**

- Requires HPC + AI/ML

# Connecting data with theory



The tool for **comprehensive validation** of cutting-edge models with multi-messenger data

→ **does a model capture the essential physics?**

**Heavy-ion event generator**

Framework for modelling all aspects of heavy-ion collisions

**Statistical toolkit**

Bayesian inference for data-theory comparison

$$P(\theta | \text{data}) = \frac{P(\text{data} | \theta)P(\theta)}{P(\text{data})}$$

**Modular** - substitute in different models

**Interdisciplinary**


(experimentalists, theorists, statisticians)

**Computing intensive calculations**

- Requires HPC + AI/ML

## State-of-the-art:

Parameter constraints with hadron and jet spectra measurements



jet transport coefficient

$$\hat{q} \sim \frac{\langle k_T \rangle}{L}$$

Steps towards **precise, quantitative** characterisation of the QGP... but **models incomplete**

# Connecting data with theory


## Next steps:

Next generation of Bayesian inference model constraints

- Roy Lemmon and myself joined JETSCAPE in 2023 to explore running next generation of calculations on UK facilities
- This year → obtained 11.8M CPU hours on CSD3 via IRIS  
→ next-gen calculations are primarily being run *here*


## Calibration

Bayesian inference to constrain 'bulk' QGP properties using 3D viscous hydrodynamics



## Model discrimination



Bayesian inference to understand how these properties (+ *different models of QGP structure*) affect jet propagation





understand jet-QGP interactions → Constrain microscopic QGP structure

# Summary

- With heavy-ion collisions, we study the deconfined state of QCD matter called **the quark-gluon plasma** - **the hottest material** that can be studied in a lab
- **Experimental upgrades** of ALICE plus **theoretical developments** means a *precision* understanding of its properties is on the horizon
- By measuring the *shockwaves* and *scatters* inside the quark-gluon plasma, we have a **unique opportunity** to uncover the microscopic dynamics of this state of matter in the near future




Rutherford, 1911






ALICE, 2020's (?)

# Backup

# Many-body QCD and the quark-gluon plasma (QGP)




## Temperature



How do fundamental degrees of freedom determine a material's emergent properties?

# Probing many-body QCD in heavy-ion collisions




We create a QGP then measure its remnants

- **How can it be investigated and its physical properties determined?**

# WP2: measuring a precise scattering probe


## How do we reconstruct charm hadrons?



Branching ratio  $\sim 6\%$

Lifetime  $c\tau \sim 60 \mu\text{m}$

C. Bartels thesis



→ Reconstruct charged particle tracks


→ Particle Identification

→ Charm hadron candidate reconstruction and selection

# WP2: measuring a precise scattering probe


## How do we reconstruct charm hadrons?

$\Lambda_c^+ \rightarrow pK^-\pi^+$   
 Branching ratio  $\sim 6\%$   
 Lifetime  $c\tau \sim 60 \mu\text{m}$




- Reconstruct charged particle tracks
- Particle Identification
- Charm hadron candidate reconstruction and selection

First ALICE paper to use Machine Learning for classification task




Signal extraction via invariant mass analysis



# WP2: measuring a precise scattering probe

## How do we reconstruct charm hadrons?




→ Reconstruct charged particle tracks

→ Particle Identification

→ Charm hadron candidate reconstruction and selection


## Charm baryon measurements probe how hadrons form

ALICE, Phys. Lett. B 839 (2023) 137796  
 Phys. Rev. Lett. 127 (2021) 202301  
 Phys. Rev. C 104 (2021) 054905  
 JHEP 04 (2018) 108






# WP3: connecting data with theory

- **Multi-messenger physics** - many measurements sensitive to same underlying physics
- Eventual aim - **Model discrimination via Bayesian inference** taking into account complete set of data




# Key challenge - TPC space charge distortions

- Drifting ions cause significant local charge which distorts driving electron path in ALICE Time Projection Chamber
- Correction maps correct distortions from up to 10cm to <1mm
- Now close to Run 2 performance




Alex Schmah,  
Quark Matter 2025

# Run 3 performance




ALI-PERF-564335



- Factor  $\sim 2$  improvement in track spatial (impact parameter) resolution in Run 3 w.r.t. Run 2

# Run 3 data




**pp:**


- 2 billion (Run1+2)
- 5700 billion (2022-2024)

\*for central barrel only (MB)  
 \*\*expected in 2025

## Goals for 2025-2026




- 47 pb<sup>-1</sup> expected
- 3 pb<sup>-1</sup> low field




- ~2.5 nb<sup>-1</sup> expected



- 2.6 nb<sup>-1</sup> expected



- 1 day + maybe one day of Ne-Ne



- ~150 nb<sup>-1</sup> expected

# Charm-tagged jets



## D<sup>0</sup>-tagged jets

**Nuclear modification factor**

$R_{AA} < 1$  - suppression



$R_{AA} = 1$  - no suppression

$$R_{AA} = \frac{1}{\langle N_{coll} \rangle} \frac{dN_{AA}/dp_T}{dN_{pp}/dp_T}$$




# Next steps: constrain interplay between wake and jets

Jet axis difference sensitive to response of QGP to jet




# Heavy-flavour correlations


- Heavy-flavour quarks produced at start of collision - provide unique way to probe the full system evolution
- ‘Diffusion’ as quark traverses medium, expected to partially thermalise depending on pT
- Correlation measurements one of most promising ‘new’ ways to constrain QGP transport properties - measuring ‘kicks’ to heavy-quark will allow for more precise probe of microscopic structure



1305.3823



Many possibilities...




# Future upgrades at the LHC

- Current ALICE setup scheduled to run until the end of Run 4 (2032), LHC scheduled to run with heavy-ions until 2041

## Longer term LHC schedule


In January 2022, the schedule was updated with long shutdown 3 (LS3) to start in 2026 and to last for 3 years. HL-LHC operations now foreseen out to end 2041.




# Inner tracker upgrade 'ITS3'

## Replacing the ITS2 inner barrel for Run 4 (2029-2032)

- bent, wafer-scale CMOS (MAPS) sensors
  - Extremely low material budget 0.02-0.04%  $X_0$
  - Homogeneous material distribution
- x2 improvement in pointing resolution, large improvement in tracking efficiency at low  $p_T$





Significant improvements in measurements of charm+beauty hadrons, plus discovery potential for exotic nuclei



# Forward Calorimeter 'FOCAL'

- Forward ( $3.4 < \eta < 5.8$ ) EM calorimeter + hadronic calorimeter to be installed for Run 4
- Exciting low-x hadron physics program, complimentary to EIC




# The ultimate heavy-ion experiment - ALICE3


ALICE3 LOI: CERN-LHCC-2022-009

## The next-generation heavy-ion experiment for LHC Run 5 and 6 (2035 onwards)

- Novel and innovative detector concept
  - Compact and lightweight all-silicon tracker
  - Retractable vertex detector
  - Extensive particle identification
  - Large acceptance
  - Superconducting magnet system
  - Continuous read-out and online processing



Ultimate performance in terms of acceptance, interaction rate and tracking precision



Precision beauty, charm correlations, exotic charm states and nuclei, dileptons, collectivity...

# Background subtraction methods for jet analyses

- **Low- $p_T$  measurements** are where ALICE is unique with respect to other LHC experiments
  - Broadening effects expected (and seen!) to be largest at low  $p_T$

- **ALICE developed tools for background at the level of *ensemble-averaged distributions***

- Data-driven
- No fragmentation bias
- Remove ALL background allowing to push down to low  $p_T$

See QM23:

N Gruenwald *EPJ Web Conf.* 296 (2024) 11005

J. Norman *EPJ Web Conf.* 296 (2024) 11006


- **Also ML approaches to background estimation**

ALICE: arXiv:2303.00592

H. Bossi, [CERN-EP seminar](#)


STAR: Phys. Rev. C 96, 024905 (2017)  
+ ALICE preliminary QM23

## Mixed event



ALICE: JHEP 09 (2015) 170

## Hadron-trigger



**Important to build on in Run 3 and beyond!**