

Event generators for collider physics

Andy Buckley

University of Glasgow

RAL Advanced Graduate Lectures, 14 June 2021



THE ROYAL
SOCIETY



University
of Glasgow




Introduction

Introduction

“Evgen” is simulation of the fundamental particle-collision process

- ▶ Ubiquitous in HEP, from experiment design to interpretation
- ▶ I'll focus on fully differential “shower+hadronisation” gens (SHGs): those that make “realistic” events that can be fed to a detector simulation
- ▶ These are serious theory tools \Rightarrow the link between “hardcore theory” and experiment. Precision (and CPU cost) have rocketed in the last decade
- ▶ But often treated as *black boxes*. . . rarely a good idea. A little extra understanding can go a long way, so let's see what we can do!




Early excuse: I'm more a user than developer. **Apologies in advance to real experts!**

Introduction

“Evgen” is simulation of the fundamental particle-collision process

- ▶ Ubiquitous in HEP, from experiment design to interpretation
- ▶ I'll focus on fully differential “shower+hadronisation” gens (SHGs): those that make “realistic” events that can be fed to a detector simulation
- ▶ These are serious theory tools \Rightarrow the link between “hardcore theory” and experiment. Precision (and CPU cost) have rocketed in the last decade
- ▶ But often treated as *black boxes*. . . rarely a good idea. A little extra understanding can go a long way, so let's see what we can do!



Early excuse: I'm more a user than developer. **Apologies in advance to real experts!**

What is an event generator?

- ▶ “Event generator” is an overloaded phrase: many pheno people refer to partonic MC integrators (e.g. MCFM, TOP++, NLOJET++, ...) as event generators. And “MC” includes Geant4!
- ▶ For experiment purposes a real EG produces *exclusive* events
 - Realistic final particle multiplicities & composition, cf. real data
 - Fortunately HEP final-states *really* can be described in full detail
- ▶ **Correlations** are not easily fakeable, e.g. from sampling data distributions: *microscopic models* produce best and most richly structured phenomena
 - i.e. event generators are based on fundamental QFT
 - but *approximately*: can't explicitly calculate full-multiplicity processes
- ▶ Since QCD is the strongest force, QCD effects usually dominate MC physics

Shower & hadronisation event generators (SHGs)

What's an SHG good for? Depends who you ask!

- ▶ **Experimentalists:** design of colliders, detectors & analyses, background estimation, signal estimation, pile-up estimation, unfolding... ~**everything!**
- ▶ **Theory/pheno:** dressing parton level calculations to make them more realistic ("easily" include effects that aren't the focus of the study e.g. decays or UE); constraining BSM models by "recasting" experimental data
- ▶ **Generator authors:** understanding (how to work with) QCD – both perturbative and non-perturbative; enabling both the above



SHGs often take in partonic events via *LHE format* (though this is evolving), and output full particle-level events via *HepMC format(s)*


A selective list

- ▶ **Partonic subprocess generators (used as SHG input via LHE):**
 - **Multi-leg LO:** MadGraph5, Sherpa, AlpGen
 - **NLO (+ multileg):** POWHEG-BOX, Sherpa, MadGraph5-aMC@NLO, MCFM
 - **Specialist processes:** HEJ, Prophecy4f, WHiZard, Protos, HEJ
- ▶ **Main general-purpose event gens:**
 - **C++:** Sherpa, Herwig 7, Pythia 8
 - **FORTRAN:** FHerwig and FPythia
- ▶ **Afterburners:**
 - EvtGen, Photos, Tauola, Jimmy
- ▶ **Specialist all-in-one:**
 - **Min bias & air showers:** PHOJET, EPOS, QGSJET, SYBILL
 - **Heavy ion:** HIJING, HYDJET, Starlight, Angantyr

Evgen in experiment data-processing

Typical experimental use of generators is to feed their output into a detector sim, e.g. based on Geant 4.


Then apply the same reconstruction + analysis as for data:



The generator bit of this chain was long considered “free” – few programs with few modes, and CPU/memory requirements *much* less than detector geometry + B -field stepping + material interaction + secondaries. **Not true these days!**

Generator capabilities, complexity & CPU demands greatly increased.

Anatomy of an SHG




ME
Shower
Hadronisation
Decays

ISR
UE
PDFs
BSM,
diffraction,
 $\tau, \gamma, B \dots$

This diag ignores UE and ISR

Anatomy of an SHG

Or, alternatively...




(by Frank Krauss)

This one is *everywhere*...

Anatomy of an SHG


Or...




(by Stefan Prestel)

Matrix elements

SHG step-by-step



SHG step-by-step



Matrix elements

Cross-sections for a scattering subprocess $ab \rightarrow n$ computed in collinear factorization:

$$\begin{aligned}\sigma &= \sum_{a,b} \int_0^1 dx_a dx_b \int f_a^{h_1}(x_a, \mu_F) f_b^{h_2}(x_b, \mu_F) d\hat{\sigma}_{ab \rightarrow n}(\mu_F, \mu_R) \\ &= \sum_{a,b} \int_0^1 dx_a dx_b \int d\Phi_n f_a^{h_1}(x_a, \mu_F) f_b^{h_2}(x_b, \mu_F) \\ &\quad \times \frac{1}{2\hat{s}} |\mathcal{M}_{ab \rightarrow n}|^2(\Phi_n; \mu_F, \mu_R),\end{aligned}$$

This is the core of all event generation: a combined *integral* of PDFs and matrix element over phase space in $x_{a,b}$ and Φ .

The “MC” comes in because the integral is done by Monte Carlo sampling in $4(n-1)$ phase space dimensions: error reduces as $\sqrt{\text{samples}}$, rather than degrading with N_{dim} .

ME integration & generation

Event generation is preceded by an integration phase to evaluate total cross-section and map the $d\hat{\sigma}$ structure in ME phase-space

Naïve sampling doesn't get far: easy to spend all CPU on *atypical points* which are dwarfed by a lucky strike on the typical set (if you're lucky!) **NLO $V+ \geq 2j$ sampling = LHC #1 MC performance challenge**

A better strategy:

- ▶ Jacobian-transform phase space to **remove divergent structures**;
- ▶ But there are *many* characteristic divergences in matrix elements
⇒ **multi-channel integration**: one per divergence, cf. MG5
- ▶ Use standard sampling techniques, or **adaptive sampling algorithms** — even ML

Integration can take *months*! Massively parallel computations becoming normal. **Typically save integration results as “gridpack” files for batched event generation.**

(Far) beyond LO

We are long past the point where SHGs only handle lowest-order partonic subprocesses (sometimes enhanced with LO *ME corrections* for the first parton shower emission).

Extra partonic emissions at tree-level increase the final state multiplicity and change the event kinematics directly. Automated by LO *merging and matching schemes* like MLM, CKKW(-L), etc.

Going beyond tree-level is more involved. An NLO cross-section has 3 parts:

$$d\sigma^{\text{NLO}} = d\tilde{\Phi}_n \left[\mathcal{B}(\tilde{\Phi}_n) + \alpha_s \mathcal{V}(\tilde{\Phi}_n) \right] + d\tilde{\Phi}_{n+1} \alpha_s \mathcal{R}(\tilde{\Phi}_{n+1})$$

But infrared divergences occur in both the Real emission and Virtual correction parts – i.e. in different Φ dimensionalities.

Bloch–Nordsieck / KLN theorems: for infra-red-safe observables, these divergences must cancel. cf. ME squaring

(Far) beyond LO

Subtraction: use **universal splitting kernel** \mathcal{S} which encodes real emission divergence structure so $\mathcal{R} - \mathcal{B} \otimes \mathcal{S}$ is finite \rightarrow computable:

$$\begin{aligned} \sigma^{\text{NLO}} = & \int_n d\tilde{\Phi}_n^{(4)} \mathcal{B} + \alpha_s \int_{n+1} d\tilde{\Phi}_{n+1}^{(4)} \left[\mathcal{R} - \mathcal{B} \otimes \mathcal{S} \right] \\ & + \alpha_s \int_n d\tilde{\Phi}_n^{(D)} \left[\tilde{\mathcal{V}} + \mathcal{B} \otimes \int_1 d\Phi_1^{(D)} \mathcal{S} \right], \end{aligned}$$

Many NLO ME calculators, but only a few automated ones.
aMC@NLO and Sherpa fully automated; POWHEG-BOX is a framework to assist manual implementation.


Virtual terms from dedicated calculators, e.g.

BlackHat/OpenLoops/NJETS/GoSAM via BLHA interface.

Technically solved: processes like $W + 5 \text{ jets}$ or **fully decayed $t\bar{t}$ and single-top** are possible... *if you can spare the integration time!*

Biased event generation and weights



- ▶ For physics purposes, we want a flat distribution of event statistics across observables
- ▶ But many distributions fall fast: if we wait for an unbiased generator to produce a TeV-scale jet, we need to make as many events as the LHC does! *At sim-reco level...*
- ▶ Neat trick: bias the sampling to produce events not from a physical distribution but from a modified one, e.g. $\hat{p}_T^4 \hat{\sigma}(\Phi)$
- ▶ Experiments usually (also) create piece-wise “sliced” samples with matched min and max cuts on $2 \rightarrow 2$ subprocess \hat{p}_T



Distributions like this are hard
to make without biasing


Biased event generation and weights

- ▶ For physics purposes, we want a flat distribution of event statistics across observables
- ▶ But many distributions fall fast: if we wait for an unbiased generator to produce a TeV-scale jet, we need to make as many events as the LHC does! *At sim-reco level...*
- ▶ Neat trick: bias the sampling to produce events not from a physical distribution but from a modified one, e.g. $\hat{p}_T^4 \hat{\sigma}(\Phi)$
- ▶ Experiments usually (also) create piece-wise “sliced” samples with matched min and max cuts on $2 \rightarrow 2$ subprocess \hat{p}_T




Parton showers


SHG step-by-step



SHG step-by-step



SHG step-by-step



Splitting functions and resummation

Some parton–jet or parton–hadron duality: can compare *fixed-order partonic events* to data for **IR-safe** observables.

Soft and collinear phase-space, $1 \text{ GeV} < p_T \lesssim 20 \text{ GeV}$, gets large *resummation corrections* from multiple QCD emissions.

Analytic resummation is process-specific and mathematically dense, especially beyond leading-log. But factorizes in the collinear limit: cross-section for process σ_0 with parton i to be accompanied by a collinear parton j with mom fraction z :

$$d\sigma \approx \sigma_0 \sum_{\text{partons}, i} \frac{\alpha_s}{2\pi} \frac{d\theta^2}{\theta^2} dz P_{ji}(z, \phi) d\phi$$

where θ is the angle between i and j and

$$\begin{aligned} P_{qq}(z) &= C_F \frac{1+z^2}{1-z}, & P_{gq}(z) &= C_F \frac{1+(1-z)^2}{z}, \\ P_{gg}(z) &= C_A \frac{z^4+1+(1-z)^4}{z(1-z)}, & P_{qg}(z) &= T_R(z^2 + (1-z)^2) \end{aligned}$$

These are the spin-averaged QCD collinear *splitting functions* – or *DGLAP kernels*. **Note divergences as $\theta, z \rightarrow 0$: dominates emissions.**

Parton showers

Parton showers are Markov chain algorithms based on the QCD splitting functions \rightarrow process-independent, approximate resummation.
Creates parton multiplicities \sim real-event hadron multiplicities

From splitting functions can calculate *probability of no emission* between scales Q and q (setting an IR cutoff for resolvability & perturbativity):

$$\Delta_i(Q^2, q^2) = \exp \left\{ - \sum_j \int_{q^2}^{Q^2} \frac{dk^2}{k^2} \frac{\alpha_s}{2\pi} \int_{q^2/k^2}^{1-q^2/k^2} dz P_{ji}(z) \right\}$$

The famous *Sudakov form factor*. Can be inverted to generate a random parton emission with physical k distribution from a random number

Split into *initial-state* (ISR) and *final-state* (FSR) showers in SHGs:

- 1 ISR: generate high- p_T extra emissions on the incoming parton legs back to the proton (using *backward evolution*);
- 2 FSR: dress all final-state partons with a forward evolution down to the QCD perturbative cutoff Λ_{QCD}

More about showers

- ▶ In practice a more complex form is used, with running α_s (and carefully chosen running scale), spin effects, quark masses, etc.
- ▶ Any evolution variable $k^2 \propto \theta^2$ is permitted in the collinear limit and will resum the divergence.
 - But some are better than others: colour coherence (cf. Chudakov effect) effects suppress emissions outside the previous emission cone.
 - Quantum effect reproduced by θ -ordering and p_T -ordering, but not virtuality. All modern generators enforce colour coherence.



More about showers

- ▶ Initial state shower adds complication
 - Emissions modifying initial partons need to change the PDF x and flavour used \Rightarrow ISR
 - Sudakovs include a PDF ratio term $\frac{x/z f_j(x/z, k^2)}{x f_i(x, k^2)}$ to \sim cancel up the chain
 - Forward-evolving from the hadron to find a consistent hard process configuration would be hopelessly inefficient \Rightarrow backward evolution.
- ▶ Actually, 1 \rightarrow 2 showers have problems:
 - Can't have finite relative p_T and real, on-shell partons since violates Lorentz symmetry \Rightarrow *reshuffling*
 - Much modern activity uses 2 \rightarrow 3 dipole showers and higher variants – also for NLO subtraction compatibility: CSS, MatchBox, Vincia, ...






Forward evolution of spacelike shower, but more fun

Shower observables

ISR: extra jets, jet distributions,
 $Z p_T$, gaps

NB. distinctly not collinear!






Shower observables

FSR: jet shapes, jet masses

i.e. adding structure to the parton = jet duality

As jet scale increases, jets become increasingly collimated



ME–shower interfacing: “matching”

Really not enough space to do this topic justice: **huge** developments in last 15 years

Issues are almost always because of **double counting** when the shower is used: an n -leg ME with parton shower *contains* the $n + 1, \dots$ terms. To improve on the Born+shower approximation, need to remove overlap.

For LO multi-leg: MLM and CKKW schemes both designed to replace the collinear shower splitting functions with proper matrix elements in the relevant (hard) phase space

Phase space slicing definitions took 10 years to iterate to better control. Introduces *merging scales*, which need to be chosen to minimise observable sensitivity: not “fire and forget” generation

Matching at NLO

Natural to go beyond tree-level matching: completely consistent NLO includes the Born, one-loop, and one-real emission amplitudes and all their interferences, *and* smoothly matches the real emissions between shower and ME.

Main benefit: NLO *scale stability*. Normalisation could always be taken from partonic highest-tech integrators, but (1-emission) shapes now also stable without disrupting formal accuracy of parton shower



⇒ **rough rule of thumb: shapes from real, normalisation from virtual**

- ▶ **Addressed first by MC@NLO:** extension of fixed-order subtraction to use shower-specific splitting functions: **process-specific**, **~ 10% negative weights**.
- ▶ **POWHEG method later:** “NLO matrix element correction”. Closer to all-positive weights, and **shower-independent**. Convenience ⇒ large uptake.

State of the art: NLO matching + merging



Now very impressive situation: automatic generation & merging of many NLO and LO multi-leg + shower in MEPS@NLO (Sherpa) and FxFx (MG5-aMC@NLO)

Bookkeeping *tour de force!* And at huge CPU cost in *unweighting*





Increasingly NLO also combined with EW corrections. NNLO is on the cards. Which features are worth the CPU?

Higher-order observables




Higher-order observables




Non-perturbative stuff
(that we wish wasn't there...)


SHG step-by-step




SHG step-by-step



SHG step-by-step




SHG step-by-step



Hadronisation

- ▶ At scales below shower cutoff $Q_0 \sim \mathcal{O}(\Lambda_{\text{QCD}})$, confinement means that physics is **non-perturbative**. **Source of most tuning params**
- ▶ Observe limited transverse momenta and Q^2 -independent energy fractions: most quantum number flow done by the shower fragmentation, so hadronisation can be \sim localised
- ▶ Two main modern hadronisation models: *Lund string and cluster*



Hadronisation models

▶ Lund string (Pythia):

- Inspired by *linear scaling of QCD potential* at large distances
- Break colour strings to produce new quark pairs; gluons form **kinks** in strings
- Lorentz invariance and LR-symmetry give Lorentz invariant *Lund symm frag function*
- Kinematics well-described, but flavour – esp. baryons – not natural

▶ Cluster hadronisation (Herwig, Sherpa):


- *Colour preconfinement*, seen in colour-connected neighbour parton mass spectrum
- Non-pert $g \rightarrow q\bar{q}$, then cluster colour singlets: requires finite gluon *constituent mass*
- Clusters treated as *meson resonances*

▶ Both models (except Sherpa) also contain *colour reconnection* heuristics

▶ **Tuning!**

Underlying event




HERA data show inclusive jet cross-section rising strongly with energy due to low- x PDFs (esp. gluon)
 \Rightarrow unitarity violation



- ▶ Eikonal models interpret the bottom-up to top-down σ ratio as mean number of *multiple-parton interactions* (MPI) \Rightarrow **sample Poisson to make n pQCD subprocesses**
- ▶ Hadron impact parameter $\sim 1/Q \Rightarrow$ transverse overlap also important
- ▶ **Low hard-process scale $Q \Rightarrow$ low overlap & low n :**
“minimum bias” cf. pile-up
- ▶ **High hard-process scale $Q \Rightarrow$ total overlap & high n :**
pedestal effect \rightarrow “underlying event”
- ▶ **Extra details:** \hat{p}_T cutoff/screening, proton overlap form factor, colour reconnection. **Tuning!**


Underlying event

HERA data show inclusive jet cross-section rising strongly with energy due to low- x PDFs (esp. gluon)
 \Rightarrow unitarity violation



MC tuning


- ▶ Freedom to describe data with generator models, via the *ad hoc* and beyond-fixed-order components: MPI, hadronisation, \sim showers
- ▶ **Need to be careful!** A pragmatic trick at LO may backfire spectacularly when a better ME is added. **Knowing the limits of a generator configuration is important**
- ▶ A global view is crucial: one number/distribution can always be overtuned at cost of others
- ▶ Rivet & Professor/Apprentice tools used to build tunes & *eigentunes*



Conclusions

Example SHG event

(Part of) a single-top HepMC event graph:



Ouch! How to safely *use* this??

Caveats on event record interpretation

- ▶ The SHG generator (or LHE) event record is often called “truth” – a dangerous phrase.
- ▶ We’re doing quantum mechanics: there is no unambiguous truth!
⇒ event records are half-physics, half-debug-info... and zero indication of amplitude interference
- ▶ It gets worse: kinematic frames aren’t defined (until the final-state) & momentum isn’t necessarily conserved at vertices!
- ▶ **BEWARE!!**



Caveats on event record interpretation

- ▶ That said, like all good myths, there is a core of truth to the widespread physical interpretation of event records
- ▶ And sometimes precision EW or PDF theorists will request correction to partonic level rather than forward-folding of their calculations, e.g. “Born-level Z ”.

NB. expts don't have to say “yes”!

- ▶ First **think about the physics** – e.g. is there a real distinction between hard photons and shower photons? Good discipline/introspection anyway!
- ▶ And first try to do what you need directly from the physical hadrons etc. See Rivet [↗](#) & ATLAS PUB note on safe truth observables [↗](#)

Apply brain!



Summary

- ▶ Event generators are super-, super-important for LHC physics
- ▶ And demands are only increasing: we demand processes and levels of data description (and predictivity) that would have been laughed at 10 years ago
- ▶ Both experiment and theory owe a great deal to the few phenomenologists who've provided us with these codes
- ▶ SHGs are based on a core of perturbative QCD (& EW) at increasingly sophisticated order. And $\sim \exp$ increasing CPU cost...
- ▶ Wrapped with perturbative iterated parton showers \Rightarrow resum logs & generate a good approximation to the high-multiplicity process we wanted. And pheno models for the stuff we don't understand *ab initio*
- ▶ **Follow-up material:** MCnet review [arXiv:1101.2599](#) · [MCnet](#) · “QCD & Collider Physics” – Ellis, Stirling, Webber · “QCD” – Dissertori, Knowles, Schmelling · MCnet summer schools: 2018 [↗](#) 2019 [↗](#)

Some hands-on exercises

Being able to run event generators yourself (rather than just using experiment samples) is a big advantage.

- ▶ Get Pythia 8 and generate QCD dijet events with ME \hat{p}_T cutoff of 50 GeV. Analyse with Rivet's MC_JETS analysis. What does it look like if you change the cutoff? How about the shower α_s ?
- ▶ Get the POWHEG-BOX heavy-flavour (hvv) process and generate $t\bar{t}$ events. Shower with Pythia and compare to top-pair production in LO Pythia using also the MC_TTBAR analysis: differences in normalisation? shapes?
- ▶ Get MG5 and generate a $p p \rightarrow z \rightarrow \mu^+ \mu^-$ process with Pythia showering. Use `add process` to merge this with +1, 2 jets MEs. Analyse with Rivet or MadAnalysis5. What changes?
- ▶ Generate a BSM process, e.g. SUSY gluino pair production in Pythia's built-in processes (you will need an SLHA file, e.g. from an experiment publication), or a FeynRules UFO model like **SMEFTsim** [↗](#) in MG5

The MCnet **hepstore** [↗](#) Docker images may be useful

More QCD and MC!

50TH INTERNATIONAL SYMPOSIUM ON MULTIPARTICLE DYNAMICS

DATE

12-16 July 2021

Online only: indico.cern.ch/e/ismd2021

TOPICS

- Hadron structure
- Forward and diffractive physics
- Collectivity and multiple scattering
- Jets and QCD at high scales
- Hadron spectroscopy
- High-temperature QCD
- Hadronic issues in heavy-flavour physics
- Cosmic ray and astrophysics



ISMD2021

