

Single-Particle Linear Dynamics

Problem Set 3

A. Wolski

April 2022

1. The *normalised* strength of a normal multipole component of a magnetic field is often specified by the “ k -value”, defined by:

$$k_n = \frac{q}{P_0} \frac{1}{n!} \left. \frac{\partial^n B_y}{\partial x^n} \right|_{x=y=0} \quad (1)$$

where q is the charge on the reference particle, P_0 is the reference momentum, and the derivative is evaluated at $x = y = 0$. Similarly, the *normalised* strength of a skew multipole component of a magnetic field is specified by a value k_{ns} , where:

$$k_{ns} = \frac{q}{P_0} \frac{1}{n!} \left. \frac{\partial^n B_x}{\partial x^n} \right|_{x=y=0} \quad (2)$$

Find the relationships between the values of k_n and k_{ns} for a multipole field, and the values of the coefficients a_n and b_n in the multipole expansion:

$$B_y + iB_x = \sum_{n=1}^{\infty} (b_n + ia_n) \left(\frac{x + iy}{r_0} \right)^{n-1} \quad (3)$$

(Note: the value of n in one representation may be different from the value of n in the other representation!)

2. A particle of charge q moves in a uniform vertical magnetic field of strength B_0 .
 - (a) Write down the Hamiltonian for the motion of the charged particle expanded to second order in the usual accelerator (normalised) variables and using path length s as the independent variable. Assume that the curvature h of the reference trajectory is equal to the normalised dipole field strength k_0 :

$$k_0 = \frac{q}{P_0} B_0 \quad (4)$$

where P_0 is the reference momentum.

- (b) Write down the equations of motion (Hamilton's equations) for the charged particle moving in this field.
- (c) Find the solutions to the equations of motion in the case that the vertical momentum is zero. Hence find the period of horizontal oscillations around the reference trajectory for a particle initially close to the reference trajectory. Express the period of the motion in terms of the circumference of the reference trajectory, and explain your answer physically.