

Extended $p_{3/2}$ Neutron Orbital and the $N = 32$ Shell Closure in ^{52}Ca

Monday, 31 July 2023 14:30 (25 minutes)

The $^{52}\text{Ca}(p,pn)^{51}\text{Ca}$ reaction was measured in inverse kinematics during the SEASTAR3 experimental campaign at the Radioactive Isotope Beam Factory (RIBF). The proton-induced quasi-free neutron knock-out reaction was performed at ~ 230 MeV/nucleon using MINOS, a 150-mm long liquid hydrogen target and the MINOS TPC, combined with prompt γ spectroscopy. Inclusive and exclusive cross sections to bound states of ^{51}Ca were evaluated, as well as the momentum distribution corresponding to the removal of $1f_{7/2}$ and $2p_{3/2}$ neutrons were measured. The cross sections, interpreted within the distorted-wave impulse approximation reaction framework, are consistent with a shell closure at the neutron number $N = 32$, found as strong as at $N = 28$ and $N = 34$ in Ca isotopes. The analysis of the momentum distributions leads to a difference of the root-mean-square radii of the neutron $1f_{7/2}$ and $2p_{3/2}$ orbitals of $0.61(23)$ fm, in agreement with the modified-shell-model prediction of 0.7 fm suggesting that the large root-mean-square radius of the $2p_{3/2}$ orbital in neutron-rich Ca isotopes is responsible for the unexpected linear increase of the charge radius with the neutron number.

Primary author: ENCIU, Madalina (INSTITUT FÜR KERNPHYSIK DER TECHNISCHEN UNIVERSITÄT DARMSTADT)

Co-authors: LIU, Hongna (Beijing Normal University); OBERTELLI, Alexandre (TU Darmstadt)

Presenter: ENCIU, Madalina (INSTITUT FÜR KERNPHYSIK DER TECHNISCHEN UNIVERSITÄT DARMSTADT)

Session Classification: Shell migration at the neutron numbers $N=32,34$ around Ca